

Analyses of multiplicity distributions by means of the Modified Negative Binomial Distribution and its KNO scaling function

Takeshi Osada, Noriaki Nakajima¹, Minoru Biyajima²
and Naomichi Suzuki³

Department of Physics, Tohoku University, Sendai 980-77, Japan

¹RCNP, Osaka University, Ibaragi 567, Japan

²Department of Physics, Shinshu University, Matsumoto 390, Japan

³Matsusho Gakuen Junior College, Matsumoto 390-12, Japan

February 1, 2008

Abstract

We analyze various data of multiplicity distributions by means of the Modified Negative Binomial Distribution (MNBD) and its KNO scaling function, since this MNBD explains the oscillating behavior of the cumulant moment observed in e^+e^- annihilations, $h-h$ collisions and $e-p$ collisions. In the present analyses, we find that the MNBD (discrete distributions) describes the data of charged particles in e^+e^- annihilations much better than the Negative Binomial Distribution (NBD).

To investigate stochastic property of the MNBD, we derive the KNO scaling function from the discrete distribution by using a straightforward method and the Poisson transform. It is a new KNO function expressed by the Laguerre polynomials. In analyses of the data by using the KNO scaling function, we find that the MNBD describes the data better than the gamma function.

Thus, it can be said that the MNBD is one of useful formulas as well as NBD.

1 Introduction

Recently it has been found that cumulant moments of the multiplicity distributions in both e^+e^- annihilations and hadronic collisions show prominent oscillatory behavior when plotted as a function of their order q [1]. In Ref.[1] this behavior is attributed to the QCD-type of branching processes apparently taking place in those reactions. However, in Refs.[2, 3] we have shown that the same behavior of the moments emerges essentially from the modified negative binomial distribution (MNBD) (which actually describes the

data of *negatively charged particles* much better than the negative binomial distribution (NBD) [4, 5, 6]). This distribution can be derived from the pure birth (PB) process with an initial condition given by the binomial distribution[7]¹.

In this paper, first of all, we analyze various experimental data of *charged particles* [9]-[20] including e - p collisions[21] by the MNBD and the NBD, to elucidate why the MNBD describes the data better than the latter in e^+e^- annihilations [2, 3].

Second, we derive the KNO scaling function of the MNBD both by the straightforward method (i.e., proceeding to the limit of large multiplicities n and large average multiplicities $\langle n \rangle$ while keeping the scaling variable $z = n/\langle n \rangle$ finite and fixed) and by the Poisson transform. Using this KNO scaling function we analyze the observed multiplicity distributions in e^+e^- annihilations, h - h collisions and e - p collisions.

Finally the concluding remarks are given. The generalized MNBD is discussed in Appendix.

2 Analyses of data by discrete distributions

It is known that the MNBD is obtained from the following equation governing the PB stochastic process[4, 5, 6, 7],

$$dP(n, t)/dt = -\lambda n P(n, t) + \lambda(n-1)P(n-1, t) \quad (1)$$

with an initial condition

$$P(n, t=0) = {}_N C_n \alpha^n (1-\alpha)^{N-n}, \quad (0 \leq \alpha \leq 1). \quad (2)$$

Here λ is a birth rate of particles. The parameter t describes the evolution of the branching processes. The generating function (GF) of the distribution at $t=0$ is given by

$$\begin{aligned} \Pi(u; t=0) &\equiv \sum_{n=0}^{\infty} P(n, t=0) \times u^n \\ &= \sum_{n=0}^{\infty} {}_N C_n \alpha^n (1-\alpha)^{N-n} u^n \\ &= [\alpha u + 1 - \alpha]^N, \end{aligned} \quad (3)$$

The GF for the MNBD at $t = \int_0^T dt = T$ is given as Refs.[2, 3, 7],

$$\Pi(u; T) \equiv \sum_{n=0}^{\infty} P(n, T) \times u^n$$

¹ It is interesting to mention here that it comes also from the concept of purely bosonic sources as presented recently in Ref.[8].

$$= \left[\frac{1 - r_1(u-1)}{1 - r_2(u-1)} \right]^N, \quad (4)$$

where

$$r_1 = -\alpha(1+p) + p,$$

$$r_2 = p \quad (\geq 0),$$

$$p = \exp(\lambda T) - 1,$$

and N (an integer) corresponds to the number of possible excited hadrons at the initial stage. The term λT is evaluated by $\lambda T = \ln(1 + r_2)$. The MNBD is given by the GF, Eq.(4), as

$$\begin{aligned} P(0) &= \Pi(0) = \left[\frac{1+r_1}{1+r_2} \right]^N, \\ P(n) &= \frac{1}{n!} \frac{\partial^n \Pi(u)}{\partial u^n} = \frac{1}{n!} \left(\frac{r_1}{r_2} \right)^N \left(\frac{r_2}{1+r_2} \right)^n \sum_{j=1}^N {}^N C_j \frac{\Gamma(n+j)}{\Gamma(j)} \left(\frac{r_2-r_1}{r_1} \right)^j \frac{1}{(1+r_2)^j}, \end{aligned} \quad (5)$$

$$\begin{aligned} r_1 &= \frac{1}{2} \left(C_2 - 1 - \frac{1}{N} \right) \langle n \rangle - \frac{1}{2}, \\ r_2 &= \frac{1}{2} \left(C_2 - 1 + \frac{1}{N} \right) \langle n \rangle - \frac{1}{2}. \end{aligned}$$

Equation (5) is applied to various experimental data[9]-[21].

For the sake of comparison with results obtained by Eq.(5), we also use the negative binomial distribution (NBD) given by

$$P(n) = \frac{\Gamma(n+k)}{\Gamma(n+1)\Gamma(k)} \left(\frac{k}{\langle n \rangle} \right)^k \left(1 + \frac{k}{\langle n \rangle} \right)^{-(n+k)}, \quad (6)$$

where $k > 0$. The GF of the NBD is given by

$$\Pi(u) = \left[1 - \frac{\langle n \rangle}{k} (u-1) \right]^{-k}. \quad (7)$$

The GF, Eq.(4), of the MNBD reduces to that of the NBD, if $r_1 = 0$.

Both the MNBD and the NBD are applied to analyses of the experimental data in e^+e^- annihilations[9]-[16], $h-h$ collisions[17]-[20] and $e-p$ collisions[21]. Since there are maximum values of multiplicity observed, n_{max} , we introduce a possible bound for N and truncate the multiplicity distribution $P(n)$ at n_{max} and renormalize it as follows,

$$\begin{aligned} N &\leq n_{max} \quad , \\ C \times \sum_{n=0}^{n_{max}} P(n) &= 1 \quad . \end{aligned}$$

In analyses in Ref.[5, 6], authors used the following treatments; 1) In the first case, the data $\langle n \rangle$ is used. r_1 and N are free. Only a constraint, $r_2 = r_1 + \langle n \rangle / N$, is used. 2) In the second case, $\langle n \rangle$, r_1 and N are free. $r_2 = r_1 + \langle n \rangle / N$ is used.

On the other hand we use C_2 and $\langle n \rangle$ as inputs. Only N is free. Our results by means of Eqs.(5) and (6) are given in Table I and some typical results of fitting are also given in Figs.1(a) and (b). It is found that in e^+e^- annihilations, the minimum values of χ^2 's obtained by fitting Eq.(5) to the data are much smaller than those of the NBD. This fact corresponds to our previous work in which we have found that the $H_q \equiv K_q / F_q$ moments obtained from the MNBD are much better than those of the NBD in e^+e^- annihilation[2,3]: However in h - h collisions they are almost equivalent.

Because the data for e - p collisions by H1 Collaborations[21] have recently reported, we calculate H_q moments by the MNBD and the NBD. Two results of comparisons are shown in Figs.2(a) and (b) for energies $\sqrt{s} = 115$ -150 GeV and $\sqrt{s} = 150$ -185 GeV, respectively. It is found that the results of the MNBD and the NBD are almost equivalent as observed in h - h collisions[3].

3 The KNO scaling function

3.1 The straightforward method

In order to know stochastic property of the MNBD, we consider the KNO scaling function of Eq.(5). Traditionally the KNO scaling function is derived from the multiplicity distribution $P(n)$ multiplied by the corresponding mean multiplicity $\langle n \rangle$ by going to the large multiplicity n and large mean multiplicity $\langle n \rangle$ limit while keeping their ratio, $z = \lim_{n, \langle n \rangle \rightarrow \infty} n / \langle n \rangle$ fixed. In our case, starting from Eq.(5) we arrive at the following function

$$\begin{aligned} \Psi(z) &\equiv \lim_{n, \langle n \rangle \rightarrow \infty} \langle n \rangle P(n) \\ &= \left(\frac{r'_1}{r'_2} \right)^N e^{-\frac{\langle n \rangle}{r'_2} z} \sum_{j=1}^N {}_N C_j \frac{1}{\Gamma(j)} \left(\frac{r'_2 - r'_1}{r'_1} \right)^j \left(\frac{\langle n \rangle}{r'_2} \right)^j z^{j-1}. \end{aligned} \quad (8)$$

The parameters r'_1 and r'_2 in Eq.(8) are given by

$$\begin{aligned} r'_1 &= \frac{1}{2} \left(C_2 - 1 - \frac{1}{N} \right) \langle n \rangle, \\ r'_2 &= \frac{1}{2} \left(C_2 - 1 + \frac{1}{N} \right) \langle n \rangle, \end{aligned} \quad (9)$$

which are slightly different from r_1, r_2 in Eq.(5), because $\langle n \rangle \gg 1$. It should be noticed that the normalization of Eq.(8) differs from the unity,

$$\int_0^\infty \Psi(z) dz = 1 - \left(\frac{r'_1}{r'_2} \right)^N, \quad (10)$$

where the second term corresponds to the term $\langle n \rangle P(0)$ in Eq.(5).

3.2 The Poisson transform

The KNO scaling function $\Psi(z, t)$ is related to the multiplicity distribution function $P(n, t)$ by the Poisson transform[22]

$$P(n, t) \xrightleftharpoons[\text{Poisson trans.}]{\text{inverse Poisson trans.}} \Psi(z, t) .$$

In this approach we obtain that

$$P(n, t) = \int_0^\infty \frac{(\alpha\omega)^n}{n!} e^{-\alpha\omega} \Psi\left(\frac{\omega}{\langle n \rangle / \alpha}, t\right) \frac{d\omega}{\langle n \rangle / \alpha}, \quad (11)$$

$$\begin{aligned} \Psi\left(\frac{\omega}{\langle n \rangle / \alpha}, t\right) &= \frac{1}{2\pi} e^{\alpha\omega} \frac{\langle n \rangle}{\alpha} \int_{-\infty}^\infty e^{-ix\omega} \sum_{n=0}^\infty \left(\frac{ix}{\alpha}\right)^n P(n, t) dx \\ &= \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} e^{sz} \Pi\left(1 - \frac{s}{\langle n \rangle}, t\right) ds, \end{aligned} \quad (12)$$

where $\Pi(u, t)$ is the generating function of $P(n, t)$. These equations hold also at $t = 0$ (the stationary function).

Using the generating function Eq.(4), the KNO scaling function $\Psi(z)$ is given by the following inverse Laplace transform

$$\Psi(z) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} e^{sz} \left(\frac{r_1}{r_2}\right)^N \sum_{j=0}^N {}_N C_j \left(\frac{r_2 - r_1 \langle n \rangle}{r_1 r_2}\right)^j \left(s + \frac{\langle n \rangle}{r_2}\right)^{-j} ds. \quad (13)$$

Then we arrive at the KNO scaling function for the MNBD,

$$\begin{aligned} \Psi(z) &= \left(\frac{r'_1}{r'_2}\right)^N \delta(z - \epsilon) + \left(\frac{r'_1}{r'_2}\right)^N e^{-\frac{\langle n \rangle}{r'_2} z} \sum_{j=1}^N {}_N C_j \frac{1}{\Gamma(j)} \left(\frac{r'_2 - r'_1}{r'_1}\right)^j \left(\frac{\langle n \rangle}{r'_2}\right)^j z^{j-1} \\ &= \left(\frac{r'_1}{r'_2}\right)^N \delta(z - \epsilon) + \left(\frac{r'_1}{r'_2}\right)^N \frac{r'_2 - r'_1 \langle n \rangle}{r'_2 r'_1} e^{-\frac{\langle n \rangle}{r'_2} z} L_{N-1}^{(1)}\left(-\frac{r'_2 - r'_1 \langle n \rangle}{r'_2 r'_1} z\right), \end{aligned} \quad (14)$$

where $L_n^{(\alpha)}(x)$ is the associated Laguerre's polynomial. In Eq.(14) the first term corresponds to the constant term in Eq.(13) (, or $\langle n \rangle P(0)$). In the numerical calculations the first term is very small because $(r'_1/r'_2)^N$ where $(N > 1)$. Therefore Eqs.(7) and (13) are almost equivalent in numerical analyses. The generalized MNBD is discussed in Appendix.

The C_q moment of the MNBD is given by

$$\begin{aligned}
C_q &\equiv \int_0^\infty dz z^q \Psi(z) \\
&= \left(\frac{r'_1}{r'_2}\right)^N e^{-\frac{\langle n \rangle}{r'_2} z} \sum_{j=1}^N {}^N C_j \frac{1}{\Gamma(j)} \left(\frac{r'_2 - r'_1}{r'_1}\right)^j \left(\frac{\langle n \rangle}{r'_2}\right)^j z^{j-1+q} \\
&= \left(\frac{r'_1}{r'_2}\right)^N \left(\frac{r'_2}{\langle n \rangle}\right)^q \sum_{j=1}^N {}^N C_j \frac{\Gamma(j+q)}{\Gamma(j)} \left(\frac{r'_2 - r'_1}{r'_1}\right)^j
\end{aligned} \tag{15}$$

We also analyze the data by the gamma function which is the KNO scaling function obtained from Eq.(6);

$$\Psi^{NBD}(z) = \frac{k^k}{\Gamma(k)} e^{-kz} z^{k-1}. \tag{16}$$

3.3 Analysis of the experimental data

We investigate the applicability of the MNBD as presented in its KNO form (i.e., using Eq.(14)) to the description of the observed multiplicity distributions in e^+e^- annihilations[9]-[16], $h-h$ collisions[17, 20] and $e-p$ collisions[21]. In actual analysis the values of the r'_1 and r'_2 are determined by using the experimental data of C_2 and $\langle n \rangle$ as mentioned before. Table II shows obtained parameters N in Eq.(14) and minimum values of χ^2 's of fitting to the experimental data. See Figs.3(a) and (b). We show here some typical results of fittings. The results using Eq.(16) are also shown there. As is seen from Table II, the minimum values of χ^2 's for the MNBD fitting are a little smaller than those for the NBD fitting in low energies e^+e^- annihilations and $h-h$ collisions (below $\sqrt{s} \sim 50$ GeV). For $e-p$ collisions the minimum values of χ^2 's are also smaller than those of the NBD up to energy $\sqrt{s}=220$ GeV.

On the other hand, in high-energy (LEP and SPS energies for the e^+e^- annihilations and $h-h$ collisions, respectively), values of the minimum χ^2 's by the MNBD fitting are almost equivalent to those of the NBD fitting.

4 Concluding remarks

Through analyses of the data by discrete distributions, we find that the MNBD describes the data in e^+e^- annihilation much better than the NBD. See Table I. We find that $N = 7 \sim 13$ except for the data by HRS and TASSO Collabs. On the other hand, in [5, 6] $N = 7 \sim 8$ for negatively charged particles is obtained, provided that their treatments for parameters are used. In other words, we have to pay our attention to the determination of parameters contained in the formulas.

Second the H_q moments of e - p collisions by H1 Collab. are analyzed by two distributions. The results are almost the same as h - h collisions. The χ^2 's in Table I are slightly better than those in Refs.[5, 6], when our treatment for parameters is used.

Third, the KNO scaling function of the MNBD is obtained. As far as our knowledge for the KNO scaling functions[24] is correct, we have not known the KNO scaling function expressed by the Laguerre polynomials. It is applied to the analyses of the observed multiplicity distribution in e^+e^- annihilations, h - h collisions, and e - p collisions. The data are also analyzed by the gamma distribution. As is seen from Table II, the MNBD describes the experimental data better (for e^+e^- annihilations) than or as well as (for h - h and e - p collisions) the NBD.

In conclusion, it can be stressed that the MNBD is useful as well as the NBD. See also Ref.[25].

What we have found in the present analyses suggests us the following: In e^+e^- annihilation, the stochastic pure birth process with the binomial distribution at $t=0$ is useful, because the finite number (corresponding to $N = 7 \sim 13$) of the excited hadrons (or the pair creation of quarks) is probably expected. In other words, the binomial distribution as the initial condition is more realistic than other condition ($\delta_{n,k}$) in the stochastic approach.

To know stochastic property of the MNBD, we have considered the generalized MNBD. In its concrete application, we have known that the discrete distribution of Eq.(A5) can not explain the oscillating behaviors of the cumulant moment observed in e^+e^- annihilations and in hadronic collisions *much better* than the MNBD, Eq.(5). For example, χ^2 's values for discrete analysis of data at $\sqrt{s}=900$ GeV are as follows: $(N, k, \chi^2)=(1, 2.22, 65.2)$, $(2, 1.33, 63.0)$ and $(3, 0.06, 60.8)$. These results suggest us that it is difficult to determine the best combination of parameters, because of three parameters. It seems to be necessary more carefulness and skillfulness than the use of the MNBD. Thus the generalized MNBD is given in Appendix. However, we are expecting that Eq.(A5) will become useful in analyses of data of some reactions at higher energies, since it has both stochastic characteristics of the MNBD and the NBD.

Acknowledgments: One of the authors (M. B.) is partially supported by the Grant-in Aid for Scientific Research from the Ministry of Education, Science, Sports and Culture (No. 06640383) and (No. 09440103). T.O. would like to thank those who supported him at the Department of Physics of Shinshu University. N. S. thanks for the financial support by Matsusho Gakuen Junior College. We are grateful to G. Wilk for his reading the manuscript.

Appendix:

In order to know the stochastic structure of the MNBD in detail, we discuss the following point: The solution obtained from the branching equation of the pure birth process with the immigration under the initial condition of the binomial distribution[3, 7] is one of

the extensions of both the MNBD and the NBD. In this case the stochastic equation (1) should be changed as follows:

$$dP(n, t)/dt = -\lambda n P(n, t) + \lambda(n-1)P(n-1, t) - \lambda_0 P(n, t) + \lambda_0 P(n-1, t), \quad (A1)$$

where λ_0 is an immigration rate. Its generating function is given as

$$\Pi(u) = \sum_{n=0}^{\infty} P(n)u^n = [1 - \tilde{r}_1(u-1)]^N [1 - \tilde{r}_2(u-1)]^{-k-N}. \quad (A2)$$

where

$$\tilde{r}_1 = \frac{\langle n \rangle}{k} \left\{ 1 - \sqrt{\frac{k+N}{N} \left[-k \left(C_2 - 1 - \frac{1}{\langle n \rangle} \right) + 1 \right]} \right\} \quad (A3)$$

$$\tilde{r}_2 = \frac{N\tilde{r}_1 + \langle n \rangle}{k+N}, \quad (A4)$$

and $k = \lambda_0/\lambda$. The MNBD is obtained by neglecting the power k in $\Pi(u)$. The generating function of the NBD is given by neglecting the power N in Eq.(A2). The physical meaning of the immigration term k may be interpreted as a possible contribution from constituent quarks and gluons. Using Eq.(4), we have directly the KNO scaling function for Eq.(A2)

$$\begin{aligned} \Psi(z) &= \left(\frac{\tilde{r}_1}{\tilde{r}_2} \right)^N e^{-\frac{\langle n \rangle}{\tilde{r}_2} z} \sum_{j=0}^N {}_N C_j \frac{1}{\Gamma(k+j)} \left(\frac{\tilde{r}_2 - \tilde{r}_1}{\tilde{r}_1} \right)^j \left(\frac{\langle n \rangle}{\tilde{r}_2} \right)^{k+j} z^{k+j-1} \\ &= \frac{\Gamma(N+1)}{\Gamma(N+k)} \left(\frac{\tilde{r}_1}{\tilde{r}_2} \right)^N \left(\frac{\langle n \rangle}{\tilde{r}_2} \right)^k z^{k-1} e^{-\frac{\langle n \rangle}{\tilde{r}_2} z} L_N^{(k-1)} \left(-\frac{\tilde{r}_2 - \tilde{r}_1}{\tilde{r}_2} \frac{\langle n \rangle}{\tilde{r}_1} z \right). \end{aligned} \quad (A5)$$

Here the term $1/\langle n \rangle$ in the parenthesis in the squared root of Eq.(A3) is neglected because $\langle n \rangle \rightarrow \infty$. It also noticed that all factors $\langle n \rangle$'s in Eq.(A5) are canceled out. This function becomes the KNO scaling function of the MNBD Eq.(14) when $k \rightarrow 0$, and reduces to the gamma distribution Eq.(16) if $N = 0$.

References

- [1] I. M. Dremin and V. A. Nechitailo, JETP Lett. **58** (1993) 881; I. M. Dremin and R. Hwa, Phys. Rev. **D49** (1994) 5805; I. M. Dremin, Phys. Lett. **B341** (1994) 95.
- [2] N. Suzuki, M. Biyajima and N. Nakajima, Phys. Rev. **D53** (1996) 3582 and **D54** (1996) 3653.
- [3] N. Nakajima, M. Biyajima and N. Suzuki, Phys. Rev. **D54** (1996) 4333.
- [4] P. V. Chliapnikov and O. G. Tchikilev, Phys. Lett. **B242** (1990) 275.

- [5] P. V. Chliapnikov, O. G. Tchikilev, and V. A. Uvarov, Phys. Lett. **B352** (1995) 461.
- [6] O. G. Tchikilev, Phys. Lett. **B382** (1996) 296, *ibid* **388**(1996),848.
- [7] N. Suzuki, M. Biyajima and G. Wilk, Phys. Lett. **B268** (1991) 447.
- [8] M. Biyajima, N. Suzuki, G. Wilk and Z. Włodarczyk. Phys. Lett. **B386** (1996) 279.
- [9] HRS Collab., M. Derrick et al., Phys. Rev. **D34** (1986) 3304.
- [10] TASSO Collab., W. Braunschweig et al., Z. Phys. **C45** (1989) 193.
- [11] AMY Collab., H. W. Zheng et al., Phys. Rev. **D42** (1990) 737.
- [12] ALEPH Collab., D. Decamp et al., Phys. Lett. **B273** (1991) 181.
- [13] ALEPH Collab., D. Buskulic et al., Z. Phys. **C69** (1995) 15.
- [14] DELPHI Collab., P. Abreu et al., Z. Phys. **C52** (1991) 271.
- [15] OPAL Collab., P. D. Acton et al., Z. Phys. **C53** (1992) 539.
- [16] OPAL Collab., P. D. Acton et al., CERN-PPE/96-47.
- [17] A. Firestone et al., Phys. Rev. **D10**(1974) 2080.
- [18] E743 Collab. R. Ammer et al., Phys. Lett. **B178**(1986) 124.
- [19] ISR Collab., A. Breakstone et al., Phys. Rev. **D30** (1984) 528.
- [20] UA5 Collab., G. J. Alner et al., Phys. Rep. **154** (1987) 247; UA5 Collab., R. E. Ansorge et al., Z. Phys. **C43** (1989) 357.
- [21] H1 Collab., S. Aid et al., DESY preprint 96-160.
- [22] M. Biyajima, Prog. Theor. Phys. **69**(1983) 966; *ibid* **70**(1983) 1468A.
- [23] A. Giovannini, S. Lupia and R. Ugoccini, Phys. Lett. **B374**, (1996) 231. F. Becattini, A. Giovannini and S. Lupia, Z. Phys. **C72** (1996), 491.
- [24] S. Saleh, ‘Photoelectron Statistics’, (Springer-Verlag, Berlin, 1998).
- [25] R. Szwed, G. Wrochna, and A.K. Wróblewski, Mod. Phys. Lett. **A6**, (1991), 245.

Table captions

Table I

The minimum values of χ^2 's for fitting to the experimental data using the MNBD and the NBD, respectively. The data of $\langle n \rangle$ and C_2 are used. We also show N giving the minimum values of χ^2 's in Eq.(6). In analysis of the data by UA5 Collab ($\sqrt{s}=900$ GeV), we find a minimum χ^2 at $N=3$ giving $P(n) < 0$ at small integer n . We discard this and adopt $N = 4$ giving $P(n) > 0$ for all n . In analysis of the data by DELPHI Collab. $\langle n \rangle = 21.26$ and $C_2 = 1.0908$ are calculated from the data.

The symbol * denotes the minimum value of χ^2 's between 2 columns. χ^2 's in Ref.[6] are obtained by means of the statistical errors only.

Table II

The same as Table I but for fitting of the KNO scaling function obtained from the MNBD and the NBD, respectively. The data of $\langle n \rangle$ and C_2 are used. In analysis of the data by OPAL ($\sqrt{s}=133$ GeV) and UA5 ($\sqrt{s}=900$ GeV) Collabs, we find minimum χ^2 's at $N = 8$ and $N = 3$ giving $P(n) < 0$ in small z , respectively. We discard these and adopt $N = 10$ and $N = 4$ giving $P(n) > 0$ for all n for the data by OPAL and UA5 Collabs, respectively.

The symbol * denotes the minimum value of χ^2 's between 2 columns.

Figure captions

[Fig. 1] Results obtained by discrete distributions.

(a) The result of fitting by Eqs.(5) and (7) to the data observed by TASSO Collab [10]. ($\sqrt{s}=34.8$ GeV.)

(b) The same as (a) but for the data observed by OPAL Collab [16]. ($\sqrt{s}=91.2$ GeV.)

[Fig.2(a) and (b)] Analyses of H_q moment in $e-p$ collisions by H1 Collaboration for energies (a) $\sqrt{s}=115-150$ GeV and (b) $\sqrt{s}=150-185$ GeV.

[Fig. 3(a) and (b)] Results obtained by the KNO scaling functions.

The same as Figs.1(a) and (b) but for fittings Eqs.(14) and (16).

Exp.	[GeV]	n_{max}	MNBD		NBD
TASSO	14	26	[N=9]	*9.34/10	25.0/11
TASSO	22	28	[N=11]	*2.14/11	10.1/12
HRS	29	28	[N=20]	*6.24/11	8.20/12
TASSO	34.8	36	[N=13]	*11.8/15	39.8/16
TASSO	43.6	38	[N=38]	*7.42/16	*7.40/17
AMY	50.	38	[N=10]	*2.75/16	14.6/17
AMY	57.	40	[N=12]	*9.84/17	51.8/18
AMY	60.	40	[N=12]	*5.79/17	11.4/18
SLD	91.2	50	[N=13]	*27.1/21	154./22
L3	91.2	50	[N=9]	*15.0/20	30.6/21
ALEPH '91	91.2	48	[N=11]	*9.07/22	12.4/23
ALEPH '95	91.2	54	[N= 9]	*3.42/25	8.97/25
DELPHI	91.2	52	[N=10]	*14.2/22	185./23
OPAL	91.2	54	[N=9]	*4.03/24	47.5/25
OPAL	133.	54	[N=7]	*4.57/21	39.6/22
FNAL	23.9	28	[N=28]	*27.1/11	34.1/12
ISR	30.4	34	[N=5]	*24.6/14	36.8/15
FNAL	38.8	32	[N=32]	*55.7/13	81.0/14
ISR	44.5	38	[N=5]	*5.27/16	16.6/17
ISR	52.6	42	[N=7]	*6.41/18	*6.96/19
ISR	62.2	40	[N=40]	*24.9/17	29.2/18
UA5	200	64	[N=4]	*7.72/28	9.87/29
UA5	546	112	[N=4]	79.1/44	*76.1/45
UA5	900	122	[N=4]	84.6/51	*77.5/52
H1 $1 < \eta^* < 5$	80-115	18	[N=18]	*2.93/16	2.92/17
	Ref.[6]		[N=7]	2.70/18	
	115-150	21	[N=21]	8.52/19	*7.01/20
	Ref.[6]		[N=7]	10.6/21	
	150-185	22	[N=22]	*5.12/20	5.65/21
	Ref.[6]		[N=7]	4.60/22	
	185-220	23	[N=23]	*5.99/21	8.97/22
	Ref.[6]		[N=7]	11.3/23	

Table I

Exp.	GeV	MNBD		NBD
TASSO	14	[N=29]	*10.9/11	19.0/12
TASSO	22	[N=28]	*3.53/12	15.3/13
HRS	29	[N=28]	*15.0/12	37.4/13
TASSO	34.8	[N=35]	*8.85/16	42.7/17
TASSO	43.6	[N=38]	*14.1/17	49.9/18
AMY	50	[N=38]	*4.19/17	*4.45/18
AMY	57	[N=29]	*7.82/18	19.1/19
AMY	60	[N=19]	*5.35/18	8.71/19
SLD	91.2	[N=19]	*35.3/22	200./23
L3	91.2	[N=11]	*13.3/21	*13.4/22
ALEPH '91	91.2	[N=15]	*8.51/22	9.95/23
ALEPH '95	91.2	[N=10]	*3.29/24	*3.59/25
DELPHI	91.2	[N=12]	*17.1/23	21.0/24
OPAL	91.2	[N=11]	*4.46/25	*4.39/26
OPAL	133	[N=10]	11.2/22	*9.63/23
FNAL	23.9	[N=28]	*26.7/12	77.1/13
ISR	30.4	[N=6]	*21.0/15	23.1/16
FNAL	38.8	[N=32]	*45.7/14	131./15
ISR	44.5	[N=6]	*5.38/17	8.19/18
ISR	52.6	[N=8]	*5.94/19	33.9/20
ISR	62.2	[N=40]	*23.5/18	59.7/19
UA5	200	[N=4]	*7.70/29	*7.69/30
UA5	546	[N=4]	67.6/45	*59.5/46
UA5	900	[N=4]	78.2/52	*63.5/53
H1 $1 < \eta^* < 5$	80-115	[N=18]	*20.5/17	37.7/18
	115-150	[N=21]	*19.5/20	32.8/21
	150-185	[N=22]	*9.55/21	18.4/22
	185-220	[N=23]	*12.3/22	23.3/23

Table II





